

About New House Farm





Over-wintered stubbles

“Big 3” Resource:
Winter Seed Food



Conservation Headlands

“Big 3” Resource:
Spring-summer invertebrate food



Pollen & Nectar Mix

“Big 3” Resource:
Spring-summer invertebrate food



Fallow Plots

“Big 3” Resource:
Places to nest in-field



Fallow Plots

“Big 3” Resource:
Places to nest in-field



Skylark Plots

“Big 3” Resource:
Places to nest in-field



Field Margins

“Big 3” Resource:
Spring-summer invertebrate food



Low input spring cereals

“Big 3” Resource:
Spring-summer invertebrate food



Field Corner Management

“Big 3” Resource:
Spring-summer invertebrate food



Wild Bird Seed Mix

“Big 3” Resource:
Winter Seed Food



Corn Bunting Mix

“Big 3” Resource:
Places to nest/Winter
Seed Food



Brassica Fodder Crops

“Big 3” Resource:
Winter Seed Food



Species Rich Grassland

“Big 3” Resource:
Spring-summer invertebrate food



Making a difference

A recent survey by one of the county bird recorders noted 300 corn buntings (beating the previous record of 225 for the area), 500 skylark, 250 yellowhammer, 100 chaffinch and 1000 linnet in the vicinity of the farm.



Species recorded on farm

Barn Owl	Garden Warbler	Lesser Whitethroat	Skylark
Blackbird	Golden Warbler	Linnet	Song Thrush
Blackcap	Goldfinch	Little Owl	Starling
Blue Tit	Great Spotted Woodpecker	Long-tailed Tit	Stock Dove
Bullfinch	Great Tit	Magpie	Stonechat
Buzzard	Green Woodpecker	Mallard	Swallow
Carrion Crow	Greenfinch	Marsh Tit	Swift
Chaffinch	Heron	Meadow Pipit	Tree Pipit
Chiffchaff	Hobby	Mistle Thrush	Wheatear
Coal Tit	House Martin	Pheasant	Whitethroat
Collared Dove	Jackdaw	Pied Wagtail	Willow Warbler
Corn Bunting	Jay	Red-legged Partridge	Wood Pigeon
Curlew	Kestrel	Redstart	Wren
Duncock	Lapwing	Robin	Yellow Wagtail
Fieldfare	Lesser black-backed gull	Rook	Yellowhammer

